

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

FURTHER MATHEMATICS

9231/12

Paper 1

May/June 2017

3 hours

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF10)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of a calculator is expected, where appropriate.

Results obtained solely from a graphic calculator, without supporting working or reasoning, will not receive credit.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **27** printed pages and **1** blank page.



BLANK PAGE

- 1 It is given that $\sum_{r=1}^n u_r = n^2(2n + 3)$, where n is a positive integer.

(i) Find $\sum_{r=n+1}^{2n} u_r$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find u_r . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2** Prove, by mathematical induction, that $5^n + 3$ is divisible by 4 for all non-negative integers n . [5]

[illegible]

3 A curve C has equation $\tan y = x$, for $x > 0$.

(i) Use implicit differentiation to show that

$$\frac{d^2y}{dx^2} = -2x \left(\frac{dy}{dx} \right)^2. \quad [3]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence find the value of $\frac{d^2y}{dx^2}$ at the point $(1, \frac{1}{4}\pi)$ on C . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 (i) Find the value of k for which the set of linear equations

$$\begin{aligned}x + 3y + kz &= 4, \\4x - 2y - 10z &= -5, \\x + y + 2z &= 1,\end{aligned}$$

has no unique solution.

[3]

This image shows a single page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

5 The matrix $\mathbf{A}$, given by

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & -2 \\ 6 & 4 & -6 \\ 6 & 5 & -7 \end{pmatrix},$$

has eigenvalues $1, -1$ and -2 .

(i) Find a set of corresponding eigenvectors.

[4]

[illegible]

- [illegible]

6 Let $I_n = \int_0^{\frac{1}{2}\pi} x^n \sin x \, dx$.

(i) Prove that, for $n \geq 2$,

$$I_n + n(n-1)I_{n-2} = n\left(\frac{1}{2}\pi\right)^{n-1}. \quad [4]$$

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

[illegible]

- 7** By finding a cubic equation whose roots are α , β and γ , solve the set of simultaneous equations

$$\alpha + \beta + \gamma = -1,$$

$$\alpha^2 + \beta^2 + \gamma^2 = 29,$$

$$\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = -1.$$

[8]

[illegible]

- 8** (i) Let $z = \cos \theta + i \sin \theta$. Show that $z - \frac{1}{z} = 2i \sin \theta$ and hence express $16 \sin^5 \theta$ in the form $\sin 5\theta + p \sin 3\theta + q \sin \theta$, where p and q are integers to be determined. [6]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

[Turn over

- 9 The curve C has equation $y = \frac{x^2 - 3x + 6}{1 - x}$.

(i) Find the equations of the asymptotes of C . [3]

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the coordinates of the turning points of C . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iii) Find the coordinates of any intersections with the coordinate axes. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iv) Sketch C . [3]

10 It is given that $x = t^{\frac{1}{2}}$, where $x > 0$ and $t > 0$, and y is a function of x .

(i) Show that $\frac{dy}{dx} = 2t^{\frac{1}{2}} \frac{dy}{dt}$ and $\frac{d^2y}{dx^2} = 2 \frac{dy}{dt} + 4t \frac{d^2y}{dt^2}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence show that the differential equation

$$\frac{d^2y}{dx^2} - \left(8x + \frac{1}{x}\right) \frac{dy}{dx} + 12x^2y = 4x^2e^{-x^2} \quad (*)$$

reduces to the differential equation

$$\frac{d^2y}{dt^2} - 4 \frac{dy}{dt} + 3y = e^{-t}. \quad [1]$$

.....

.....

.....

.....

.....

.....

.....

.....

This image shows a full page of white paper with horizontal dotted lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

11 The curve C has polar equation $r = a(1 + \sin \theta)$ for $-\pi < \theta \leq \pi$, where a is a positive constant.

(i) Sketch C .

[2]

(ii) Find the area of the region enclosed by C .

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Show that the length of the arc of C from the pole to the point furthest from the pole is given by

$$s = (\sqrt{2})a \int_{-\frac{1}{2}\pi}^{\frac{1}{2}\pi} \sqrt{1 + \sin \theta} \, d\theta. \quad [3]$$

[illegible]

[illegible]

12 Answer only **one** of the following two alternatives.

EITHER

The curve C has equation $y = \frac{1}{2}(e^x + e^{-x})$ for $0 \leq x \leq 4$.

- (i) The region R is bounded by C , the x -axis, the y -axis and the line $x = 4$. Find, in terms of e , the coordinates of the centroid of the region R . [10]

[illegible]

- (ii) Show that $\frac{ds}{dx} = \frac{1}{2}(e^x + e^{-x})$, where s denotes the arc length of C , and find the surface area generated when C is rotated through 2π radians about the x -axis. [4]

[illegible]

$$\mathbf{i} + \mathbf{j} + 3\mathbf{k}, \quad 3\mathbf{i} - \mathbf{j} + 5\mathbf{k}, \quad 3\mathbf{i} - \mathbf{j} + \mathbf{k}, \quad 5\mathbf{i} - 5\mathbf{j} + \alpha\mathbf{k},$$

(i) Show that the possible values of α are 3 and 5. [7]

[illegible]

-
-
-
-
-
-
-
- (ii) Using $\alpha = 3$, find the shortest distance of the point D from the line AC , giving your answer correct to 3 significant figures. [3]

- (iii) Using $\alpha = 3$, find the acute angle between the planes ABC and ABD , giving your answer in degrees. [4]

[illegible]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.