

**MARK SCHEME for the October/November 2009 question paper
for the guidance of teachers**

9231 FURTHER MATHEMATICS

9231/02

Paper 2, maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2009	9231	02

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol $\checkmark$ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2009	9231	02

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
CWO	Correct Working Only – often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{}$ " marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2009	9231	02

Question Number	Mark Scheme Details	Part Mark	Total
4	Find MI of rod and ring about A: $I = \frac{1}{3} 2ma^2 + 2ma^2 + ma^2$ M1 Use conservation of energy: $\frac{1}{2} I (d\theta/dt)^2 = 3mga \sin \theta$ M1 Substitute $I = (11/3) ma^2$: $a (d\theta/dt)^2 = (18/11)g \sin \theta$ A.G. A1 (3) Resolve forces on ring along rod (ignore sign of F): $F = mg \sin \theta + ma (d\theta/dt)^2$ M1 Substitute for $d\theta/dt$: $F = (29/11) mg \sin \theta$ A.G. A1 Resolve forces on ring normal to rod (ignore signs): $R = mg \cos \theta - ma d^2\theta/dt^2$ M1 Find $d^2\theta/dt^2$ by differentiating eqn above: $a d^2\theta/dt^2 = (9/11)g \cos \theta$ M1 A1 Combine to give magnitude of normal force: $R = (2/11) mg \cos \theta$ A.G. A1 (6) Relate μ to θ: $\mu = F/R = (29/2) \tan \theta$ M1 Find θ: $\theta = \tan^{-1} (2\mu/29)$ A1 (2) [11]		
5	Equate moments of AB and BC about A: $Wa \sin \alpha = Wa (\cos \alpha - 2 \sin \alpha)$ M1 A1 Evaluate α: $\tan \alpha = \frac{1}{3}$, $\alpha = 18.4^\circ$ A.G. A1 (3) Take moments about B for rod BC: $T 2a \cos 45^\circ = Wa \cos \alpha$ M1 A1 Find tension T in form kW: $T = (\cos 18.4^\circ / \sqrt{2}) W = 0.671 W$ A1 (3) <i>EITHER:</i> Resolve vertically for rod BC: $R_V = W - T \sin (45^\circ + \alpha)$ (ignore signs of all components R_V etc) $= W - T \sin 63.4^\circ$ [= 0.400 W] M1 A1 Resolve horizontally for rod BC: $R_H = T \cos (45^\circ + \alpha)$ [= 0.300 W] M1 A1 <i>OR:</i> Resolve along AB for rod BC: $R_{AB} = W \cos \alpha - T \sin 45^\circ$ (or take moments about C) [= $3W/2\sqrt{10} = 0.474 W$] (M1 A1) Resolve along BC for rod BC: $R_{BC} = W \sin \alpha - T \cos 45^\circ$ [= $-W/2\sqrt{10} = -0.158 W$] (M1 A1) Combine components, e.g. $\sqrt{(R_V^2 + R_H^2)}$: $R = 0.5 W$ M1 A1 (6) [12]		

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2009	9231	02

Question Number	Mark Scheme Details	Part Mark	Total
6	Find sample mean (2 dp required): $\bar{x} = 60.18 / 6 = 10.03$ B1 Estimate population variance to 3 sf (allow biased here: 0.00107 or 0.0327²): $s^2 = (603.6118 - 60.18^2/6) / 5 = 0.00128 \text{ or } 0.0358^2$ B1 Use consistent formula for C.I. with any t value: $\bar{x} \pm t\sqrt{(s^2/6)}$ M1√ Use of correct tabular value: $t_{5, 0.995} = 4.03[2]$ *A1 C.I. correct to 2 dp in cm (dep *A1): 10.03 ± 0.06 or [9.97, 10.09] A1 (5) Find CI for mean circumference to 1 dp in cm: $10.03\pi \pm 0.06\pi$ $= 31.5 \pm 0.2 \text{ or } [31.3, 31.7]$ B1 (1) [6]		
7	Find $P(X \geq 3/4)$: $1 - F(3/4) = 1 - 1/2[(3/4)^3 + 1]$ M1 $= 37/128 \text{ or } 0.289$ *A1 State deduction about upper quartile Q_3 (dep *A1): $Q_3 > 3/4$ B1 (3) Express cum. dist. fn. G of Y in terms of X: $G(y) = P(Y \leq y) = P(-\sqrt{y} \leq X \leq \sqrt{y})$ M1 A1 Relate to F: $= F(\sqrt{y}) - F(-\sqrt{y})$ M1 Simplify: $= 1/2[(\sqrt{y})^3 + 1] - 1/2[(-\sqrt{y})^3 + 1]$ $= y^{3/2}$ A1 State G in full (√ on previous result): $0 \ (y < 0), y^{3/2} \ (0 \leq y \leq 1), 1 \ (y > 1)$ A1√ (5) [8]		
8	State if assumption necessary with valid reason: No, since large samples (A.E.F.) B1 State another valid assumption: Samples must be random (A.E.F.) B1 State hypotheses: $H_0: \mu_c - \mu_f = 5, H_1: \mu_c - \mu_f < 5$ B1 Find z (A.E.F.): $z = (19.14 - 15.36 - 5) /$ $\sqrt{(20.54/75 + 9.84/75)}$ M1 A1 $= -1.22 / \sqrt{(2 \times 15.19/75)} = -1.92$ *A1 Use correct tabular value of z: $z_{0.95} = 1.64[5]$ *B1 Consistent conclusion (A.E.F.; A1 dep *A1, *B1): Does exceed by less than 5 kg. M1√ A1 (8) [9]		

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2009	9231	02

Question Number	Mark Scheme Details	Part Mark	Total
9	Show how value is found (allow M1 if 100 omitted): $100 {}^4C_2 0.4^2 \times 0.6^2 = 34.56$ A.G. M1 A1 State (at least) null hypothesis: $H_0: B(4, 0.6)$ fits data (A.E.F.) B1 Combine adjacent cells since exp. value < 5: $O: \begin{matrix} 14 & 27 & 49 & 10 \end{matrix}$ $E: \begin{matrix} 17.92 & 34.56 & 34.56 & 12.96 \end{matrix}$ *M1 Calculate value of χ^2 (to 2 dp ; A1 dep *M1): $\chi^2 = 9.22$ M1 A1 Compare with consistent tabular value (to 2 dp): $\chi_{3, 0.95}^2 = 7.815$ (cells combined) $\chi_{4, 0.95}^2 = 9.488$ (not combined) B1 Valid method for reaching conclusion: Reject H_0 if $\chi^2 >$ tabular value M1 Correct conclusion (A.E.F., requires correct values): $9.22 > 7.81[5]$ so distn. does not fit *A1 State valid deduction (dep *A1; allow $p \neq 0.6$): Prob. of faulty chips $\neq 0.6$ (A.E.F.) B1	(2)	
		(8)	[10]
10	Use geometric distribution [with $p = 3/20 = 0.15$] M1 Find prob. of missing in 5 shots: q^5 or $1 - p(1 + q + q^2 + q^3 + q^4)$ M1 $(q^6$ or 0.377 earns M1 A0) $= 0.85^5 = 0.444$ A1 Separate probabilities of hits: $P(Y = r) = P(2 \text{ hits in } r-1 \text{ shots}) \times$ $P(\text{hit on } r^{\text{th}} \text{ shot})$ M1 Substitute using geometric distribution: $= {}^{r-1}C_2 \times 0.15^2 \times 0.85^{r-3} \times 0.15$ M1 Replace ${}^{r-1}C_2$: A.G. $= \frac{1}{2}(r-1)(r-2) 0.15^3 \times 0.85^{r-3}$ A1 Simplify $P(Y = r+1) / P(Y = r)$: $0.85 r / (r-2)$ or $17r / 20(r-2)$ B1 Find required values of r: $17r < 20(r-2)$ M1 $r > 40/3$ or 13.3 or 13 A1 Deduce most probable value of Y: 14 ($\sqrt{\text{ on values of } r}$) A1✓	(3)	
		(3)	
		(4)	[10]

Page 9	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2009	9231	02

Question Number	Mark Scheme Details			Part Mark	Total
11 OR	State what is minimised (A.E.F.):	Sum of squares of residuals	B1		
	Show residuals on copy of diagram:	Verticals between points and line	B1	(2)	
	State answer with valid reason (A.E.F.) e.g.:	No, since points not collinear	B1	(1)	
	(i) Calculate correlation coefficient:				
		$r = (1022 \cdot 15 - 340 \times 22 \cdot 41 / 8) / \sqrt{\{(15\,500 - 340^2 / 8)(67 \cdot 65 - 22 \cdot 41^2 / 8)\}}$	M1 A1		
		$= 69 \cdot 725 / \sqrt{(1050 \times 4 \cdot 874)} = 0 \cdot 975$	*A1		
	State valid comment on diagram (A.E.F.):	Points lie close to a straight line	B1✓		
	(B0 for 'points are nearly correlated')			(4)	
	(ii) State valid reason (A.E.F.; dep *A1):	0·975 closer to 1 than 0·965	B1		
	Find coefficient b in regression line for y :	$b = (1022 \cdot 15 - 340 \times 22 \cdot 41 / 8) / (15\,500 - 340^2 / 8)$			
		$= 69 \cdot 725 / 1050 = 0 \cdot 0664$	M1 A1		
	Find equation of regression line:	$\sqrt{z} = b(v - 42 \cdot 5) + 2 \cdot 80125$			
		$= 0 \cdot 0664 v - 0 \cdot 0207[5]$			
		<i>or</i> $0 \cdot 066 v - 0 \cdot 021$	M1 A1	(5)	
	(iii) All answers A.E.F.:				
	<i>EITHER</i> : State physical significance:	Equivalent to $\sqrt{z}$ for zero speed	B1		
	<i>OR</i> : State expected relationship:	Constant force so $v^2 \propto z$	(B1)		
	Valid comment, $\sqrt{}$ on 0·021:	$0 \cdot 021$ <i>or</i> $0 \cdot 021^2 \approx 0$ as expected			
		<i>or</i> $\neq 0$ suggests error in data	B1	(2)	
					[14]