Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

0123456789

FURTHER MATHEMATICS **9231/01**

Paper 1 Further Pure Mathematics 1 **For examination from 2020**

SPECIMEN PAPER **2 hours**

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Blank pages are indicated.

1 **(a)** Given that $f(r) = \dfrac{1}{(r+1)(r+2)}$, show that

$$f(r-1) - f(r) = \frac{2}{r(r+1)(r+2)}.$$ [2]

(b) Hence find $\sum_{r=1}^{n} \dfrac{1}{r(r+1)(r+2)}$. [3]

(c) Deduce the value of $\sum_{r=1}^{\infty} \dfrac{1}{r(r+1)(r+2)}$. [1]

2 It is given that $\phi(n) = 5^n(4n + 1) - 1$, for $n = 1, 2, 3, \ldots$.

Prove, by mathematical induction, that $\phi(n)$ is divisible by 8 for every positive integer n. [7]

3 The curve C has polar equation $r = 2 + 2\cos\theta$, for $0 \leqslant \theta \leqslant \pi$.

(a) Sketch C. [3]

(b) Find the area of the region enclosed by C and the initial line. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Show that the Cartesian equation of C can be expressed as $4(x^2 + y^2) = (x^2 + y^2 - 2x)^2$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 The cubic equation

$$z^3 - z^2 - z - 5 = 0$$

has roots α, β and γ.

(a) Show that the value of $\alpha^3 + \beta^3 + \gamma^3$ is 19. [4]

(b) Find the value of $\alpha^4 + \beta^4 + \gamma^4$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find a cubic equation with roots $\alpha + 1$, $\beta + 1$ and $\gamma + 1$, giving your answer in the form

$$px^3 + qx^2 + rx + s = 0,$$

where p, q, r and s are constants to be determined. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 The matrix **A** is given by

$$\mathbf{A} = \begin{pmatrix} 5 & k \\ -3 & -4 \end{pmatrix}.$$

(a) Find the value of k for which **A** is singular. [2]

.....

.....

.....

.....

.....

.....

.....

It is now given that $k = 6$ so that $\mathbf{A} = \begin{pmatrix} 5 & 6 \\ -3 & -4 \end{pmatrix}$.

(b) Find the equations of the invariant lines, through the origin, of the transformation in the x-y plane represented by **A**. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) The triangle *DEF* in the *x*-*y* plane is transformed by **A** onto triangle *PQR*.

(i) Given that the area of triangle *DEF* is $10\,\text{cm}^2$, find the area of triangle *PQR*. [2]

.....

.....

.....

.....

.....

.....

(ii) Find the matrix which transforms triangle *PQR* onto triangle *DEF*. [2]

.....

.....

.....

.....

.....

6 The position vectors of the points A, B, C, D are

$$2\mathbf{i} + 4\mathbf{j} - 3\mathbf{k}, \quad -2\mathbf{i} + 5\mathbf{j} - 4\mathbf{k}, \quad \mathbf{i} + 4\mathbf{j} + \mathbf{k}, \quad \mathbf{i} + 5\mathbf{j} + m\mathbf{k},$$

respectively, where m is an integer. It is given that the shortest distance between the line through A and B and the line through C and D is 3.

(a) Show that the only possible value of m is 2. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the shortest distance of D from the line through A and C. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Show that the acute angle between the planes ACD and BCD is $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$. [4]

7 The curve C has equation $y = \dfrac{2x^2 - 3x - 2}{x^2 - 2x + 1}$.

(a) State the equations of the asymptotes of C. [2]

.....

.....

.....

.....

.....

(b) Show that $y \leqslant \frac{25}{12}$ at all points on C. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the coordinates of any stationary points of C. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Sketch C, stating the coordinates of any intersections of C with the coordinate axes and the asymptotes. [4]

(e) Sketch the curve with equation $y = \left|\dfrac{2x^2 - 3x - 2}{x^2 - 2x + 1}\right|$ and find the set of values of x for which $\left|\dfrac{2x^2 - 3x - 2}{x^2 - 2x + 1}\right| < 2$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....